



11239 Ventura Blvd., Suite 213
Studio City, CA 91604
Phone: (818) 505-0152
Fax: (818) 505-0398

Ryan Greene D.O.
Garrett Glapa, ACNP-BC
Abe Grosswasser D.C.

PATIENT QUESTIONS

Activity Level (Select all that apply):

- Low (Sedentary)
- Moderate (Walk/jog/workout infrequently)
- Average (Walk/jog/workout 1 to 3 times per week)
- High (Walk/jog/workout regularly 4+ times per week)

SYMPTOMS AND CONCERNS

Select all that apply:

- | | |
|---|--|
| <input type="checkbox"/> Acne | <input type="checkbox"/> Decrease in Strength or Endurance |
| <input type="checkbox"/> Erectile Dysfunction (ED) | <input type="checkbox"/> Decrease in Work Performance |
| <input type="checkbox"/> Decreased Libido | <input type="checkbox"/> Frequent Urinary Tract Infection |
| <input type="checkbox"/> Decreased Desire | <input type="checkbox"/> Brittle Nails |
| <input type="checkbox"/> Inability To or Delayed Orgasm | <input type="checkbox"/> Thinning Eyebrows |
| <input type="checkbox"/> Weight Gain | <input type="checkbox"/> Hair Thinning |
| <input type="checkbox"/> Decreased Muscle Mass | <input type="checkbox"/> Cold Hands or Feet |
| <input type="checkbox"/> Difficulty Sleeping | <input type="checkbox"/> Mind Racing at Bedtime |
| <input type="checkbox"/> Urinary Incontinence | <input type="checkbox"/> Mood Swings |
| <input type="checkbox"/> Dry or Flaking Skin | <input type="checkbox"/> Gynecomastia |
| <input type="checkbox"/> Lack of Energy (Fatigue) | <input type="checkbox"/> Abdominal Obesity |

Male Hormone Replacement Insertion Consent

Name: _____ Date of Birth: _____

My physician/practitioner has recommended testosterone therapy delivered by a pellet inserted under my skin for treatment of symptoms I am experiencing related to low testosterone levels. The following information has been explained to me prior to receiving the recommended testosterone therapy.

OVERVIEW

Bioidentical testosterone is a form of testosterone that is biologically identical to that made in my own body. The levels of active testosterone made by my body have decreased, and therapy using these hormones may have the same or similar effect(s) on my body as my own naturally produced testosterone. The pellets are a delivery mechanism for testosterone, and bioidentical hormone replacement therapy using pellets has been used since the 1930's. There are other formulations of testosterone replacement available, and different methods can be used to deliver the therapy. The risks associated with pellet therapy are generally similar to other forms of replacement therapy using bioidentical hormones.

RISKS/COMPLICATIONS

Risks associated with pellet insertion may include: bleeding from incision site, bruising, fever, infection, pain, swelling, pellet extrusion, which may occur several weeks or months after insertion, reaction to local anesthetic and/or preservatives, allergy to adhesives from bandage(s), steri strips or other adhesive agents.

Some individuals may experience one or more of the following complications: acne, anxiety, breast or nipple tenderness or swelling, insomnia, depression, mood swings, fluid and electrolyte disturbances, headaches, increase in body hair, fluid retention or swelling, mood swings or irritability, rash, redness, itching, lack of effect (typically from lack of absorption), transient increase in cholesterol, nausea, retention of sodium, chloride and/or potassium, weight gain or weight loss, thinning hair or male pattern baldness, increased growth of prostate and prostate tumors, which may or may not lead to worsening of urinary symptoms, hypersexuality (overactive libido) or decreased libido, erectile dysfunction, painful ejaculation, ten to fifteen percent shrinkage in testicular size, and/or significant reduction in sperm production, which may lead to permanent infertility, and increased hematocrit. The latter can be diagnosed with a blood test called a complete blood count (CBC). This test should be done at least annually. Elevated HCT, which is dose and serum testosterone level dependent, may lead to increased clotting.

All types of testosterone replacement can cause a significant decrease in sperm count during use. If you are planning to start or expand your family, please talk to your provider about other non-testosterone options.

It is advisable that all males > 40 years old who are on testosterone therapy should have a PSA checked at least yearly.

CONSENT FOR TREATMENT

I agree to immediately report any adverse reactions or problems that may be related to my therapy to my physician or health care provider's office, so that it may be reported to the manufacturer. Potential complications have been explained to me, and I acknowledge that I have received and understand this information, including the possible risks and potential complications and the potential benefits. I also acknowledge that the nature of bioidentical therapy and other treatments have been explained to me, and I have had all my questions answered.

Blood tests may be necessary on several occasions during the 1st year to help with dosing, and then annually or biannually at the discretion of the prescribing practitioner.

I understand that my blood tests may reveal that my levels are not optimal, which would mean I may need a higher or lower dose in the future. Furthermore, I have not been promised or guaranteed any specific benefits from the insertion of testosterone pellets. I have read or have had this form read to me.

I accept these risks and benefits, and I consent to the insertion of testosterone pellets under my skin performed by my provider. This consent is ongoing for this and all future insertions in this facility until I am no longer a patient here, but I do understand that I can revoke my consent at any time. I have been informed that I may experience any of the complications to this procedure as described above.

I ACKNOWLEDGE THAT I HAVE RECEIVED A COPY AND UNDERSTAND THE INFORMATION ON THIS FORM

PATIENT:

Print Name: _____

Signature: _____ Date: _____

WITNESS:

Print Name: _____

Signature: _____ Date: _____